

Chapter 11 Orthogonal Functions and

正交

Fourier Series

傅立葉

complete and
orthogonal)

複習：linear algebra 關於 orthogonal (正交) basis 的介紹

在 linear algebra 當中

$$(1) \text{ inner product } (\mathbf{f}_1, \mathbf{f}_2) = \sum_n \mathbf{f}_1[n] \mathbf{f}_2[n]$$

$[1, 2, 1]$
 and $[-2, 2, -2]$
 are orthogonal
 If A is an orthogonal matrix

$$(2) \text{ orthogonal } \sum_n \mathbf{f}_1[n] \mathbf{f}_2[n] = 0$$

$$\mathbf{A}^T \mathbf{A} = \mathbf{D}$$

$$\mathbf{D}^{-1} \mathbf{A}^T \mathbf{A} = \mathbf{I}$$

(3) 若 $\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_N$ 為 complete orthogonal set,

$$\mathbf{A}^{-1} = \mathbf{D}^{-1} \mathbf{A}^T$$

inner
 product $\mathbf{f}[n] = \sum_{m=1}^N a_m \mathbf{f}_m[n]$ where $a_m = \frac{\sum_{n=1}^N \mathbf{f}[n] \mathbf{f}_m[n]}{\sum_{n=1}^N \mathbf{f}_m[n] \mathbf{f}_m[n]}$

$$\downarrow (\mathbf{f}[n], \mathbf{f}_e[n]) = \sum_{m=1}^N a_m (\mathbf{f}_m[n], \mathbf{f}_e[n])$$

$$= a_e (\mathbf{f}_e[n], \mathbf{f}_e[n])$$


例如 在只有三個 entry 的情形下

$$\mathbf{f}_1 = \begin{bmatrix} 1 \\ f_1[1] \\ f_1[2] \\ f_1[3] \end{bmatrix}$$

$$\mathbf{f}_2 = \begin{bmatrix} 0 \\ f_2[1] \\ f_2[2] \\ f_2[3] \end{bmatrix}$$

$$\mathbf{f}_3 = \begin{bmatrix} 0 \\ f_3[1] \\ f_3[2] \\ f_3[3] \end{bmatrix}$$

是一組 complete orthogonal set



問題：在 continuous 當中該如何定義 orthogonal?

Section 11.1 Orthogonal Functions

11.1.1 緝要：熟悉幾個重要定義

- (1) inner product (page 347) (7) normalize (page 354)
- (2) orthogonal (page 349) (8) complete (page 356)
- (3) orthogonal set (page 351) (9) orthogonal series expansion (page 358)
- (4) square norm (page 353) (10) generalized Fourier series (page 358)
- (5) norm (page 353) (11) weight function (page 360)
- (6) orthonormal set (page 353) ↑
With weighting functions, many definitions are changed.

學習方式：(1) 可以多和 linear algebra 當中的定義多比較
 (2) 複習三角函式的公式 (see page 363-364)

11.1.2 定義

(1) inner product on an interval $[a, b]$

$$\star_1 (f_1, f_2) = \int_a^b f_1(x) f_2(x) dx \quad (f_1, f_2 \text{ 為 real 時})$$

比較 : discrete case $(\mathbf{f}_1, \mathbf{f}_2) = \sum_n \mathbf{f}_1[n] \mathbf{f}_2[n]$

補充 : more standard definition for inner product

$$(f_1, f_2) = \int_a^b f_1(x) f_2^*(x) dx$$

\star_{1-1} with conjugation

Inner product 性質

(a) $(f_1, f_2) = (f_2, f_1)^*$

 $*$: conjugation

$$(f_1, kf_2) = k^*(f_1, f_2)$$

(b) $(k f_1, f_2) = k (f_1, f_2)$, k 為 scalar (或稱為constant)

(c) $\underline{(f, f) = 0}$ if and only if $f = 0$, $\underline{(f, f) > 0}$ if and only if $f \neq 0$,

 \star_{1-2}

(d) $(f_1 + f_2, g) = (f_1, g) + (f_2, g)$

discrete case 亦有這些性質

(2) orthogonal on an interval $[a, b]$

$$\text{def}_2 \quad (f_1, f_2) = \int_a^b f_1(x) f_2(x) dx = 0 \quad (f_1, f_2 \text{ 為 real 時})$$

$$\text{或 } (f_1, f_2) = \int_a^b f_1(x) f_2^*(x) dx = 0 \quad (\text{more standard definition})$$

$$\text{比較 : discrete case} \quad \sum_n \mathbf{f}_1[n] \mathbf{f}_2[n] = 0$$

例子: 當 $[a, b] = [-1, 1]$,

1 和 x^k (k 為奇數) 互為 orthogonal
 even odd

$$\int_{-1}^1 1 \cdot x^k dx = \frac{x^{k+1}}{k+1} \Big|_{-1}^1 = \frac{1^{k+1} - (-1)^{k+1}}{k+1} = \frac{1-1}{k+1} = 0$$

★₂₋₂

$$f(x) = f(-x)$$

$$f(x) = -f(-x)$$

350

注意：任何 even function 和任何 odd function 在 $[-b, b]$ 之間必為 orthogonal, ($a = -b$)


包括 Example 1 (text page 426) 的 x^2 和 x^3 在 $[-1, 1]$ 之間也是 orthogonal

$$\int_{-b}^b f_1(x) f_2(x) dx = 0$$

even odd

$$(x^2, x^3) = \int_{-1}^1 x^2 x^3 dx = \int_{-1}^1 x^5 dx = \frac{x^6}{6} \Big|_{-1}^1 = 0$$

$$\text{even} \cdot \text{odd} = \text{odd}$$



$$\int_{-b}^b \text{odd function } dx = 0$$

$\cos(x^2)$ $\tan(x^3)$ are orthogonal at $[-5, 5]$

$$\int_{-5}^5 \cos(x^2) \tan(x^3) dx = 0$$

(3) orthogonal set \star_3

有一組 functions $\phi_0(x), \phi_1(x), \phi_2(x), \phi_3(x), \dots$

若 $\int_a^b \phi_m(x) \phi_n^*(x) dx = 0$ for any $m \neq n$

則 $\phi_0(x), \phi_1(x), \phi_2(x), \phi_3(x), \dots$ 被稱作 orthogonal set on an interval $[a, b]$

Example 2 (text page 426)

Show that the set $\{1, \cos x, \cos 2x, \cos 3x, \dots\}$ is an orthogonal set on the interval $[-\pi, \pi]$

page 367, p = \pi

Case 1

when one of the functions is 1

$$\int_{-\pi}^{\pi} 1 \cdot \cos nx dx = \frac{1}{n} \sin nx \Big|_{-\pi}^{\pi} = 0$$

Case 2

when both the two functions are not 1

運用三角函式公式
(pages 363-364)

$$\int_{-\pi}^{\pi} \cos mx \cos nx dx = \int_{-\pi}^{\pi} \frac{1}{2} (\cos(m+n)x + \cos(m-n)x) dx$$

$$= \frac{\sin((m+n)\pi)}{2(m+n)} - \frac{\sin(-(m+n)\pi)}{2(m+n)} + \frac{\sin((m-n)\pi)}{2(m-n)} - \frac{\sin(-(m-n)\pi)}{2(m-n)} = 0$$

(4) square norm \star_4

$$\underline{\|f(x)\|^2} = \underline{(f(x), f(x))} = \int_a^b f(x) f^*(x) dx = \underline{\int_a^b |f(x)|^2 dx}$$

比較：discrete case $\sum_n \mathbf{f}[n] \mathbf{f}^*[n]$

(5) norm \star_5

$$\|f(x)\| = \sqrt{(f(x), f(x))} = \sqrt{\int_a^b f(x) f^*(x) dx} = \sqrt{\int_a^b |f(x)|^2 dx}$$

(6) orthonormal set \star_6

對一個 orthogonal set, 若更進一步的滿足

$$\underline{\int_a^b \phi_n(x) \phi_n^*(x) dx = 1} \quad \text{for all } n$$

則被稱為 orthonormal set

(7) normalize

 \star_7 norm
 $\neq 1$

normalize

norm = 1

將 norm 變為 1

 $\psi(x)$
/sat/

normalize

$$v(x) = \frac{\psi(x)}{\|\psi(x)\|}$$

normalize

注意，此時

$$\underline{(v(x), v(x))} = \left(\frac{\psi(x)}{\|\psi(x)\|}, \frac{\psi(x)}{\|\psi(x)\|} \right) = \frac{1}{\|\psi(x)\|^2} (\psi(x), \psi(x)) = \underline{1}$$

可藉由 normalization, 將 orthogonal set 變成 orthonormal set
normalize


Example 3 (text page 427)

Calculate the norms of $\{1, \cos x, \cos 2x, \cos 3x, \dots\}$

$$\|1\|^2 = \int_{-\pi}^{\pi} 1 \cdot 1 dx = x \Big|_{-\pi}^{\pi} = 2\pi$$

$$\|\cos nx\|^2 = \int_{-\pi}^{\pi} \cos nx \cos nx dx = \int_{-\pi}^{\pi} \frac{1}{2}(\cos 2nx + 1) dx \quad \begin{array}{l} \text{運用三角函式公式} \\ (\text{pages 363-364}) \end{array}$$

$$= \frac{\sin 2nx}{4n} + \frac{x}{2} \Big|_{-\pi}^{\pi} = \frac{\sin 2n\pi}{4n} + \frac{\pi}{2} - \frac{\sin(-2n\pi)}{4n} - \frac{(-\pi)}{2} = \pi$$

$$\|1\| = \sqrt{2\pi} \quad \|\cos nx\| = \sqrt{\pi}$$

$\{1, \cos x, \cos 2x, \cos 3x, \dots\}$ can be normalized as an orthonormal set

$$\left\{ \frac{1}{\sqrt{2\pi}}, \frac{\cos x}{\sqrt{\pi}}, \frac{\cos 2x}{\sqrt{\pi}}, \frac{\cos 3x}{\sqrt{\pi}}, \dots \right\}$$

(8) complete  8

若在 interval $[a, b]$ 之間，任何一個 function $f(x)$ 都可以表示成 $\phi_0(x), \phi_1(x), \phi_2(x), \phi_3(x), \dots$ 的 linear combination

$$f(x) = c_0\phi_0(x) + c_1\phi_1(x) + c_2\phi_2(x) + \dots = \sum_{n=0}^{\infty} c_n\phi_n(x)$$

則 $\phi_0(x), \phi_1(x), \phi_2(x), \phi_3(x), \dots$ 被稱作 complete

比較：在 linear algebra 當中，對 3-D vector 而言

$e_1 = [1, 0, 0], e_2 = [0, 1, 0], e_3 = [0, 0, 1]$ 為 complete

Any 3-D vector $[a, b, c]$ can be expressed as $a\mathbf{e}_1 + b\mathbf{e}_2 + c\mathbf{e}_3$

In the continuous case,

Complete: infinite number of functions are required to form a complete set.

For $\phi_0(x), \phi_1(x), \phi_2(x), \phi_3(x), \dots$

if we can find $c_0, c_1, c_2, c_3, \dots$ such that

$$f(x) - \sum_{n=0}^{\infty} c_n \phi_n(x) = 0 \quad \text{for all } x$$

then $\phi_0(x), \phi_1(x), \phi_2(x), \phi_3(x), \dots$ is a complete set.

Example: $\{1, x-x_0, (x-x_0)^2, (x-x_0)^3, \dots\}$ is a complete set if $f(x)$ is constrained to be a continuous function. *see page 275, Taylor series*

However, $\{1, x-x_0, (x-x_0)^2, (x-x_0)^3, \dots\}$ are not orthogonal.

$\{1, \cos x, \cos 2x, \cos 3x, \dots\}$ are orthogonal but not complete.
(all even)

(9)(10)

若 $\phi_0(x), \phi_1(x), \phi_2(x), \phi_3(x), \dots$ 為 complete,
可將 $f(x)$ 表示成

$$f(x) = \sum_{n=0}^{\infty} c_n \phi_n(x)$$

Question: How to find c_n ?

當 $\phi_0(x), \phi_1(x), \phi_2(x), \phi_3(x), \dots$ 不為 orthogonal, c_n 不易算

當 $\phi_0(x), \phi_1(x), \phi_2(x), \phi_3(x), \dots$ 為 orthogonal

$$f(x) = \sum_{m=0}^{\infty} c_m \phi_m(x)$$

$$\int_a^b f(x) \phi_n^*(x) dx = \sum_{m=0}^{\infty} c_m \int_a^b \phi_m(x) \phi_n^*(x) dx = c_n \int_a^b \phi_n(x) \phi_n^*(x) dx$$

$\int_a^b \phi_m(x) \phi_n(x) dx = 0$ if $m \neq n$

$$c_n = \frac{\int_a^b f(x) \phi_n^*(x) dx}{\int_a^b \phi_n(x) \phi_n^*(x) dx}$$

★9

被稱作 (9) orthogonal series expansion

★10

c_n 被稱作 (10) generalized Fourier series

當 $\phi_0(x), \phi_1(x), \phi_2(x), \phi_3(x), \dots \dots \dots$ 為 orthonormal

$$c_n = \int_a^b f(x) \phi_n^*(x) dx$$

*q-1

Examples of Complete and Orthogonal Sets in the Continuous Case:

11.1.3 Orthogonal with Weight Function

加權

(11-1) inner product with weight function

★ 11-1

$$(f_1(x), f_2(x)) = \int_a^b w(x) f_1(x) f_2^*(x) dx$$

ex: $w(x) = e^{-x^2}$

其中 $w(x)$ 被稱作 weight function

加上了 weight function 後

(11-2) orthogonal 的定義改成

★ 11-2

$$(f_m, f_n) = \int_a^b w(x) f_m(x) f_n^*(x) dx = 0 \quad \text{for } m \neq n$$

(11-3) square norm 的定義改成

★ 11-3

$$\|f(x)\|^2 = \int_a^b w(x) f(x) f^*(x) dx$$

(11-4) norm 的定義改成

$$\text{11-4} \quad \|f(x)\| = \sqrt{\int_a^b w(x) f(x) f^*(x) dx}$$

(11-5) orthonormal 的定義改成

$$\text{11-5} \quad \int_a^b w(x) f_m(x) f_n^*(x) dx = 0 \quad \text{for } m \neq n$$

$$\int_a^b w(x) f_n(x) f_n^*(x) dx = 1$$

(11-6) normalize 的算法改成

$$\text{11-6} \quad v(x) = \frac{\psi(x)}{\|\psi(x)\|} = \frac{\psi(x)}{\sqrt{\int_a^b w(x) \underline{\psi}(x) \psi^*(x) dx}}$$

(11-7) orthogonal series expansion of $f(x)$ 以及 generalize Fourier

\star_{11-7} series 的算法改成

$$f(x) = \sum_{n=0}^{\infty} c_n \phi_n(x)$$

$$c_n = \frac{\int_a^b w(x) f(x) \phi_n^*(x) dx}{\int_a^b w(x) \phi_n(x) \phi_n^*(x) dx}$$

(要複習) **☆₁₂**

$\cos(a + b)$	$\cos a \cos b - \sin a \sin b$
$\sin(a + b)$	$\sin a \cos b + \cos a \sin b$
$\cos(a - b)$	$\cos a \cos b + \sin a \sin b$
$\sin(a - b)$	$\sin a \cos b - \cos a \sin b$
$\cos a \cos b$	$[\cos(a + b) + \cos(a - b)]/2$
$\sin a \sin b$	$[\cos(a - b) - \cos(a + b)]/2$
$\sin a \cos b$	$[\sin(a + b) + \sin(a - b)]/2$

A₁₂

364

$\cos(2a)$	$\cos^2 a - \sin^2 a$ or $1 - 2\sin^2 a$ or $2\cos^2 a - 1$
$\sin(2a)$	$2\sin a \cos a$
$\cos^2 a$	$[\cos(2a) + 1]/2$
$\sin^2 a$	$[1 - \cos(2a)]/2$

11.1.5 Section 11.1 需要注意的地方

(1) Norm 和 square of norm 要分清楚

做 normalization 時，要除以 norm

(2) 熟悉三角函數的公式

(i) 記住幾個，其他的就不難推算出來

(ii) 許多公式可以由 $\cos(a) = \frac{e^{ja} + e^{-ja}}{2}$ 導出來

$$\sin(a) = \frac{e^{ja} - e^{-ja}}{2j} = \frac{je^{-ja} - je^{ja}}{2}$$

Two well-known complete and orthogonal function set

(1) Fourier series, Fourier cosine series, Fourier sine series
(Sections 11-2 and 11-3)

(2) Legendre polynomials (Section 6-4)

$$\int_{-1}^1 P_m(x) P_n(x) dx = 0 \quad \text{if } m \neq n$$

其他常用的 complete and orthogonal function set

Hermite polynomials (with weight function) (補充)

Chebyshev polynomials (with weight function) (補充)

1808

1812

1822

Section 11.2 Fourier Series

傅立葉級數



11.2.1 緬要

trigonometric functions \star , complete and orthogonal

$$\left\{ 1, \cos \frac{\pi}{p}x, \cos \frac{2\pi}{p}x, \cos \frac{3\pi}{p}x, \dots, \sin \frac{\pi}{p}x, \sin \frac{2\pi}{p}x, \sin \frac{3\pi}{p}x, \dots \right\}$$

frequency = 0
(DC term)

orthogonal set on the interval of $[-p, p]$



$$\cos \frac{n\pi}{p}x \quad \text{週期 : } \frac{2p}{n} \quad \text{頻率 : } \frac{n}{2p}$$

(2) Fourier Series

Note! $\frac{a_0}{2}$, not a_0

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi}{P} x + b_n \sin \frac{n\pi}{P} x \right)$$

$x \in [-P, P]$

$a_0 = \frac{1}{P} \int_{-P}^P f(x) dx$ $a_n = \frac{1}{P} \int_{-P}^P f(x) \cos \frac{n\pi}{P} x dx$

$b_n = \frac{1}{P} \int_{-P}^P f(x) \sin \frac{n\pi}{P} x dx$

$x \in [-P, P]$

Remember them!

page 358

from

page 358

(紅色部分特別注意，勿記錯公式)

$$\begin{aligned} \frac{a_0}{2} &= \frac{\int_{-P}^P f(x) \cdot 1 dx}{\int_{-P}^P 1 \cdot 1 dx} & a_n &= \frac{\int_{-P}^P f(x) \cos \frac{n\pi}{P} x dx}{\int_{-P}^P (\cos^2 \frac{n\pi}{P} x) dx} \\ &= \frac{\int_{-P}^P f(x) dx}{2P} & \int_{-P}^P (\cos^2 \frac{n\pi}{P} x) dx &= \int_{-P}^P \frac{\cos(\frac{2n\pi}{P} x) + 1}{2} dx \\ \therefore a_0 &= \frac{\int_{-P}^P f(x) dx}{P} & &= \frac{\sin \frac{2n\pi}{P} x}{4n\pi/P} \Big|_{-P}^P + \frac{x}{2} \Big|_{-P}^P = 0 + \frac{P}{2} - \frac{-P}{2} = P \\ && a_n &= \frac{\int_{-P}^P f(x) \cos \frac{n\pi}{P} x dx}{P} \end{aligned}$$

(3) 名詞

trigonometric function (page 372)

Fourier series (trigonometric series) (page 376)

Fourier coefficients (page 376)

partial sum (page 379)

fundamental period (page 384)

period extension (page 384)

物理意義：

Fourier Series == 對信號做頻率分析

「頻率」(frequency)是個常用字，以 Hz(每秒多少個週期)為單位

說話聲音: 100~1200 Hz

人耳可聽見的聲音: 20~20000 Hz

廣播 (AM): $5 \times 10^5 \sim 1.6 \times 10^6$ Hz

廣播 (FM): $8.8 \times 10^7 \sim 1.08 \times 10^8$ Hz

無線電視: $7.6 \times 10^7 \sim 8.8 \times 10^7$, $1.74 \times 10^8 \sim 2.16 \times 10^8$ Hz

行動通訊: 5.1×10^8 Hz ~ 2.75×10^{11} Hz

可見光: 4×10^{14} Hz ~ 8×10^{14} Hz

測量頻率的方式：
Fourier series
Fourier transform

11.2.2 Trigonometric Functions

trigonometric functions

$$\left\{ 1, \cos \frac{\pi}{p}x, \cos \frac{2\pi}{p}x, \cos \frac{3\pi}{p}x, \dots, \sin \frac{\pi}{p}x, \sin \frac{2\pi}{p}x, \sin \frac{3\pi}{p}x, \dots \right\}$$

type 1
type 2
type 3

Trigonometric functions is orthogonal on the interval of $[-p, p]$

要用 $C_3^2 + 2 = 5$ 次的inner products 來證明

(1) 1 VS. Cosine

$$\int_{-p}^p 1 \cdot \cos \frac{\pi k}{p} x dx = \frac{p}{\pi k} \sin \frac{\pi k}{p} x \Big|_{-p}^p = \frac{p}{\pi k} \sin \pi k - \frac{p}{\pi k} \sin(-\pi k) = 0 - 0 = 0$$

(2) 1 VS. Sine

$$\int_{-p}^p 1 \cdot \sin \frac{\pi k}{p} x dx = -\frac{p}{\pi k} \cos \frac{\pi k}{p} x \Big|_{-p}^p = -\frac{p}{\pi k} \cos \pi k + \frac{p}{\pi k} \cos(-\pi k) = 0$$

even
odd

(3) Cosine VS. Sine

$$\begin{aligned}
 & \int_{-p}^p \cos \frac{\pi k}{p} x \cdot \sin \frac{\pi h}{p} x dx = \int_{-p}^p \frac{1}{2} \left[\sin \frac{\pi(h+k)}{p} x - \sin \frac{\pi(h-k)}{p} x \right] dx \\
 &= \frac{p}{2\pi} \left[-\frac{1}{h+k} \cos \left(\frac{\pi(h+k)x}{p} \right) + \frac{1}{h-k} \cos \left(\frac{\pi(h-k)x}{p} \right) \right] \Big|_{-p}^p \\
 &= \frac{p}{2\pi} \left[-\frac{1}{h+k} [\cos(\pi(h+k)) - \cos(-\pi(h+k))] \right. \\
 &\quad \left. + \frac{1}{h-k} [\cos(\pi(h-k)) - \cos(-\pi(h-k))] \right] = 0 \quad (\text{when } h \neq k)
 \end{aligned}$$

when ($h = k$)

$$\int_{-p}^p \cos \frac{\pi k}{p} x \cdot \sin \frac{\pi h}{p} x dx = \int_{-p}^p \frac{1}{2} \sin \frac{2\pi k}{p} x dx = -\frac{p}{4\pi k} \cos \frac{2\pi k}{p} \Big|_{-p}^p = 0$$

(4) Cosine VS. Cosine, $k \neq h$

$$\begin{aligned}
 \int_{-p}^p \cos \frac{\pi k}{p} x \cdot \cos \frac{\pi h}{p} x dx &= \int_{-p}^p \frac{1}{2} \left[\cos \frac{\pi(h+k)}{p} x + \cos \frac{\pi(h-k)}{p} x \right] dx \\
 &= \frac{p}{2\pi} \left[\frac{1}{h+k} \sin \left(\frac{\pi(h+k)x}{p} \right) + \frac{1}{h-k} \sin \left(\frac{\pi(h-k)x}{p} \right) \right]_{-p}^p \\
 &= \frac{p}{2\pi} \left[\frac{1}{h+k} [\sin(\pi(h+k)) - \sin(-\pi(h+k))] \right. \\
 &\quad \left. + \frac{1}{h-k} [\sin(\pi(h-k)) - \sin(-\pi(h-k))] \right] = 0 \quad \text{when } h \neq k
 \end{aligned}$$

(5) Sine VS. Sine, $k \neq h$

page 363

$$\begin{aligned}
 \int_{-p}^p \sin \frac{\pi k}{p} x \cdot \sin \frac{\pi h}{p} x dx &= \int_{-p}^p \frac{1}{2} \left[\cos \frac{\pi(h-k)}{p} x - \cos \frac{\pi(h+k)}{p} x \right] dx \\
 &= \frac{p}{2\pi} \left[\frac{1}{h-k} \sin \left(\frac{\pi(h-k)x}{p} \right) - \frac{1}{h+k} \sin \left(\frac{\pi(h+k)x}{p} \right) \right]_{-p}^p = 0 \quad \text{when } h \neq k
 \end{aligned}$$

Square norms of trigonometric functions

$$\|1\|^2 = \int_{-p}^p 1 \cdot 1 dx = x \Big|_{-p}^p = 2p$$

$$\left\| \cos \frac{\pi k}{p} x \right\|^2 = \int_{-p}^p \cos^2 \frac{\pi k}{p} x dx = \frac{1}{2} \int_{-p}^p (1 + \cos \frac{2\pi k}{p} x) dx = \frac{1}{2} \left(x + \frac{p}{2\pi k} \sin \frac{2\pi k}{p} x \right) \Big|_{-p}^p = p$$

$$\left\| \sin \frac{\pi kx}{p} \right\|^2 = \int_{-p}^p \sin^2 \frac{\pi kx}{p} dx = \frac{1}{2} \int_{-p}^p (1 - \cos \frac{2\pi kx}{p}) dx = \frac{1}{2} \left(x - \frac{p}{2\pi k} \sin \frac{2\pi kx}{p} \right) \Big|_{-p}^p = p$$

11.2.3 Fourier Series



The Fourier series is the orthogonal series expansion (see page 358) by trigonometric functions

(Fourier series又被稱作 trigonometric series)

The Fourier Series of a function $f(x)$ defined on the interval $[-p, p]$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi}{p} x + b_n \sin \frac{n\pi}{p} x \right)$$

式2


$$\left\{ \begin{array}{l} a_0 = \frac{1}{p} \int_{-p}^p f(x) dx \\ a_n = \frac{1}{p} \int_{-p}^p f(x) \cos \frac{n\pi}{p} x dx \\ b_n = \frac{1}{p} \int_{-p}^p f(x) \sin \frac{n\pi}{p} x dx \end{array} \right.$$

a_0, a_n, b_n 被稱作 Fourier coefficients

Example 1 (text page 433)

$$P = \pi$$

$$f(x) = \begin{cases} 0 & \text{for } -\pi < x < 0 \\ \pi - x & \text{for } 0 \leq x < \pi \end{cases}$$



$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{\pi} \int_0^{\pi} (\pi - x) dx = \frac{\pi}{2}$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx = \frac{1}{\pi} \int_0^{\pi} (\pi - x) \cos nx dx$$

$$= \frac{\pi - x}{\pi} \frac{1}{n} \sin nx \Big|_0^\pi - \frac{1}{\pi} \int_0^{\pi} (-1) \frac{1}{n} \sin nx dx$$

$$= -\frac{1}{n^2 \pi} \cos nx \Big|_0^\pi = \frac{1 - \cos n\pi}{n^2 \pi} = \frac{1 - (-1)^n}{n^2 \pi}$$

$$\star_3$$

$$\int u v' = u v - \int u' v$$

$$u = \pi - x$$

$$v' = \cos nx$$

$$v = \frac{1}{n} \sin nx$$

$$\cos n\pi = (-1)^n \star_4$$

$$\int u v' = uv - \int u' v$$

378

$$u = \pi - x$$

$$v = \frac{-\cos nx}{n}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx = \frac{1}{\pi} \int_0^{\pi} (\pi - x) \sin nx dx \\
 &= -\frac{\pi - x}{\pi} \frac{1}{n} \cos nx \Big|_0^{\pi} - \frac{1}{\pi} \int_0^{\pi} (-1)(-\frac{1}{n} \cos nx) dx \\
 &= \frac{1}{n} - \frac{1}{n^2 \pi} \sin nx \Big|_0^{\pi} = \frac{1}{n}
 \end{aligned}$$

$$\boxed{
 \begin{aligned}
 f(x) &= \frac{\pi}{4} + \sum_{n=1}^{\infty} \left(\frac{1 - (-1)^n}{n^2 \pi} \cos \frac{n\pi}{p} x + \frac{1}{n} \sin \frac{n\pi}{p} x \right) \\
 &= \frac{\pi}{4} + \sum_{n=1}^{\infty} \left(\frac{1 - (-1)^n}{n^2 \pi} \cos nx + \frac{1}{n} \sin nx \right)
 \end{aligned}
 }$$

$$p = \pi$$

11.2.4 Sequence of Partial Sums


Sequence of Partial Sums

$$S_N(x) = \frac{a_0}{2} + \sum_{n=1}^N \left(a_n \cos \frac{n\pi}{p} x + b_n \sin \frac{n\pi}{p} x \right)$$

$$f_1(x) = \lim_{N \rightarrow \infty} S_N(x)$$

N 越大，越能逼近原來的 function


For Example 1



(a) $S_3(x)$



$$N = 3$$



(b) $S_8(x)$



$$N = 8$$

Fig. 11.2.3


(c) $S_{15}(x)$  $N = 15$

Fig. 11.2.3

11.2.5 Conditions for Convergence

$$\text{If } a_0 = \frac{1}{p} \int_{-p}^p f(x) dx \quad a_n = \frac{1}{p} \int_{-p}^p f(x) \cos \frac{n\pi}{p} x dx$$

$$b_n = \frac{1}{p} \int_{-p}^p f(x) \sin \frac{n\pi}{p} x dx$$

$$f_1(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi}{p} x + b_n \sin \frac{n\pi}{p} x \right)$$

$f_1(x) = f(x)$ 其實未必成立

(1) $f_1(x_0) = f(x_0)$ if $f(x)$ is continuous at x_0

$$(2) \quad f_1(x_0) = \frac{f(x_0+) + f(x_0-)}{2} \quad \text{if } f(x) \text{ is not continuous at } x_0$$

$\star 5-1$

$$f(x_0+) = \lim_{h \rightarrow 0} f(x_0 + h) \quad f(x_0-) = \lim_{h \rightarrow 0} f(x_0 - h)$$

Example 1 的例子





Fig. 11-2-1

11.2.6 Period Extension

$$f_1(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi}{p} x + b_n \sin \frac{n\pi}{p} x \right)$$

fundamental period: $2p$

在 interval $x \in [-p, p]$ 以外的地方

★ 5-2

$$f_1(x + 2p) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \left(\frac{n\pi}{p} x + 2n\pi \right) + b_n \sin \left(\frac{n\pi}{p} x + 2n\pi \right) \right)$$

$$\underline{f_1(x + 2p)} = \underline{f_1(x)} \quad (\text{period Extension})$$

- ◆ $f_1(x)$ 是個週期為 $2p$ 的函式 (這是 $f_1(x)$ 和 $f(x)$ 第二個不同的地方)

Example 1 的例子

$$p = \pi$$




Fig. 11.2.2



對一個非週期的函式，Fourier series expansion 的結果不適用
於 $x \notin [-p, p]$ 的區域

但是週期函式則可

11.2.7 Section 11.2 需要注意的地方

(1) Fourier series 的公式 (常背錯)

(a) 第一項是 $a_0/2$ ，而非 a_0

(b) 算 a_0, a_n, b_n 時，積分後別忘了除以 p

(p 是 interval width 的一半)

(2) 背熟三角函式公式

$$(3) \text{熟悉 } \int_a^b u(t)v'(t)dt = u(t)v(t)\Big|_a^b - \int_a^b u'(t)v(t)dt$$

(在計算 Fourier coefficients 會常用到，如 Example 1)

(4) 當 n 為整數時, $\cos n\pi = (-1)^n$ 習慣這種表示法

(5) 正確而言, $f(x) \approx \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi}{p} x + b_n \sin \frac{n\pi}{p} x \right)$

↑
近似於

因為當 $f_1(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi}{p} x + b_n \sin \frac{n\pi}{p} x \right)$

$f_1(x)$ 和 $f(x)$ 之間有二個不同的地方

(a) 在 discontinuous 的地方 $f_1(x_0) = [f(x_0+) + f(x_0-)]/2$

(b) $f_1(x)$ 為 periodic, $f_1(x) = f_1(x + p)$

然而, 習慣上, 還是寫成 $f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi}{p} x + b_n \sin \frac{n\pi}{p} x \right)$

數學史上最美麗的詩篇 --- 傅立葉級數

Clerk Maxwell

悲傷的傅立葉

Section 11.3 Fourier Cosine and Sine Series

11.3.1 緬要

(1)

Fourier Series

比較 page 368

$\star\star 1-1$
 $f(x)$ is even

$$b_n = \frac{1}{P} \int_{-P}^P f(x) \sin\left(\frac{n\pi}{P}x\right) dx = 0$$

$$a_n = \frac{1}{P} \int_{-P}^P f(x) \cos\left(\frac{n\pi}{P}x\right) dx = \frac{2}{P} \int_0^P f(x) \cos\left(\frac{n\pi}{P}x\right) dx$$

Fourier cosine series (或 cosine series)

不同一： $b_n = 0$

不同二： $\frac{1}{P} \int_{-P}^P \Rightarrow \frac{2}{P} \int_0^P$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{P} x$$

$$a_0 = \frac{2}{P} \int_0^P f(x) dx \quad a_n = \frac{2}{P} \int_0^P f(x) \cos \frac{n\pi}{P} x dx$$

$\star\star 1-2$
 $f(x)$ is odd

Fourier sine series (或 sine series)

不同一： $a_n = 0, a_0 = 0$

$$f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi}{P} x$$

不同二： $\frac{1}{P} \int_{-P}^P \Rightarrow \frac{2}{P} \int_0^P$

$$b_n = \frac{2}{P} \int_0^P f(x) \sin \frac{n\pi}{P} x dx$$

(2) 重要名詞 : Fourier cosine series, cosine series (page 395)

Fourier sine series, sine series (page 396)

Gibb's Phenomenon (page 399)



(3) Half-range extension: $[0, L]$

(a) cosine series: $f(x) = f(-x)$, interval is changed into $[-L, L]$, set $p = L$

(b) sine series: $f(x) = -f(-x)$, interval is changed into $[-L, L]$, set $p = L$

(c) Fourier series: (i) interval $[-p, p]$ is replaced by $[0, L]$,

(ii) p is replaced by $L/2$

(4) One of the applications: Solving particular solution

(See pages 407-412)

11.3.2 Even and Odd Functions

even function: $f(x) = f(-x)$

odd function: $f(x) = -f(-x)$


Example

$1, x^2, x^4, x^6, x^8 \dots$ are even

$x, x^3, x^5, x^7, x^9 \dots$ are odd


Cosine functions are even

$$\cos(t)$$



Sine functions are odd

$$\sin(t)$$



Several properties about even and odd functions

(a) The product of two even functions is even

$$\text{例 : } x^2 \cdot x^4 = x^6$$

(b) The product of two odd functions is even

$$\text{例 : } x \cdot x = x^2$$

(c) The product of an even function and an odd function is odd

$$\text{例 : } x \cdot x^2 = x^3$$

(d) The sum (or difference) of two even function is still even

(e) The sum (or difference) of two odd function is still odd

(f) If $f(x)$ is even, then

$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$$

\star_{2-1}



(g) If $f(x)$ is odd, then

$$\int_{-a}^a f(x) dx = 0$$

A_{2-2}

(Proof):

$$\begin{aligned} \int_{-a}^a f(x) dx &= \int_{-a}^0 f(x) dx + \int_0^a f(x) dx \\ &= - \int_a^0 f(-x_1) dx_1 + \int_0^a f(x) dx \quad (\text{令 } x_1 = -x, dx_1 = -dx) \\ &= \int_0^a f(-x_1) dx_1 + \int_0^a f(x) dx \end{aligned}$$

When $f(x) = f(-x)$

$$\int_{-a}^a f(x) dx = \int_0^a f(x_1) dx_1 + \int_0^a f(x) dx = 2 \int_0^a f(x) dx$$

When $f(x) = -f(-x)$

$$\int_{-a}^a f(x) dx = \int_0^a -f(x_1) dx_1 + \int_0^a f(x) dx = 0$$

11.3.3 Fourier Cosine and Sine Series

(1) The Fourier series of an even function on the interval $(-p, p)$ is the cosine series (或稱作 Fourier cosine series)

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{p} x$$

$$a_0 = \frac{2}{p} \int_0^p f(x) dx$$

$$a_n = \frac{2}{p} \int_0^p f(x) \cos \frac{n\pi}{p} x dx$$

和之前 Fourier series 不一樣的地方有三個

適用情形：(1) $f(x)$ is even

(2) Half range extension (page 401)

(2) The Fourier series of an odd function on the interval $(-p, p)$ is the sine series (或稱作 Fourier sine series)

$$f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi}{p} x$$

$$b_n = \frac{2}{p} \int_0^p f(x) \sin \frac{n\pi}{p} x dx$$

和之前 Fourier series 不一樣的地方有三個
(是哪三個)

適用情形：(1) $f(x)$ is odd
(2) Half range extension (page 401)

Example 1 (text page 438) $p=2$ $f(x) = -f(-x)$ odd

397

Expand $f(x) = x, -2 < x < 2$ in a Fourier series




Fig. 11.3.3

$f(x)$ is odd
 \therefore expand $f(x)$ by a Fourier sine series
 $p=2$

$$\int uv' = uv - \int u'v
u = x, v = \frac{-2}{n\pi} \cos \frac{n\pi}{2} x$$

$$b_n = \frac{2}{2} \int_0^2 x \sin \frac{n\pi}{2} x dx = -\frac{2}{n\pi} x \cos \frac{n\pi}{2} x \Big|_0^2 + \frac{2}{n\pi} \int_0^2 \cos \frac{n\pi}{2} x dx$$

$$= -\frac{2}{n\pi} 2 \cos n\pi + 0 + \frac{4}{n^2 \pi^2} \sin \frac{n\pi}{2} x \Big|_0^2 = -\frac{4}{n\pi} (-1)^n + 0 - 0 = \frac{4}{n\pi} (-1)^{n+1}$$

$$f(x) = \sum_{n=1}^{\infty} \frac{4(-1)^{n+1}}{n\pi} \sin \frac{n\pi}{2} x$$

$$\cos n\pi = (-1)^n$$

Example 2 (text page 438)

$$f(x) = \begin{cases} -1 & -\pi < x < 0 \\ 1 & 0 \leq x < \pi \end{cases}$$





Fig. 11.3.5

square wave



$$f(x) = -f(-x)$$

odd function, 使用 sine series

$$b_n = \frac{2}{\pi} \int_0^\pi 1 \cdot \sin \frac{n\pi}{\pi} x dx = -\frac{2}{\pi} \frac{\cos nx}{n} \Big|_0^\pi = \frac{2}{\pi} \frac{1 - (-1)^n}{n}$$

$$\cos n\pi = (-1)^n$$

$$f(x) = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1 - (-1)^n}{n} \sin nx$$

11.3.4 Gibbs Phenomenon

Example 2 的結果 $f(x) = \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1 - (-1)^n}{n} \sin nx$

partial sum $S_N(x) = \frac{2}{\pi} \sum_{n=1}^{N} \frac{1 - (-1)^n}{n} \sin nx$

當 N 不為無限大，在 discontinuities 附近會有 “overshooting”

“overshooting”的大小不會隨著 N 而變小

但寬度會越來越窄，越來越靠近 discontinuities 的地方

這種現象，稱作 Gibb's phenomenon







Fig. 11.3.6

11.3.5 Half Range Extension

★3

之前的例子： $f(x)$ is defined in the interval of $-p < x < p$

若問題改成

Expand $f(x)$, $0 < x < L$

$(f(x)$ 只有在 $0 < x < L$ 當中有定義
對範圍之外的 $f(x)$ 用猜測的方式)

★3-1

(a) In a cosine series, suppose that $\underline{\underline{f(x)}} = \underline{\underline{f(-x)}}$ and $\underline{\underline{f(x)}} = \underline{\underline{f(x+2L)}}$

(i) Interval: $[-L, L]$, (ii) 所有公式的 p 由 L 取代, (iii) 結果是 even

★3-2

(b) in a sine series, suppose that $f(x) = -f(-x)$ and $f(x) = f(x+2L)$

(i) Interval: $[-L, L]$, (ii) 所有公式的 p 由 L 取代, (iii) 結果是 odd

★3-3

(c) in a Fourier series, suppose that $f(x) = f(x+L)$

Note!

(i) Interval: $[0, L]$, (ii) 所有公式的 p 由 $L/2$ 取代
(和前面兩個不同)

如 Example 3 (text page 440),

$$f(x) = x^2, \quad 0 < x < L$$

(a) in a cosine series

假設 $f(x) = f(-x)$ for $-L < x < 0$, (假設 $f(x)$ 是一個 even function)
interval 變為 $(-L, L)$

原本 cosine series 公式

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{p} x$$

$$f(x) = f(x+2p) \quad x \in [-p, p]$$

$$a_0 = \frac{2}{p} \int_0^p f(x) dx$$

$$a_n = \frac{2}{p} \int_0^p f(x) \cos \frac{n\pi}{p} x dx$$

現在只不過將 p 改成 L

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{L} x$$

$$a_0 = \frac{2}{L} \int_0^L f(x) dx$$

$$a_n = \frac{2}{L} \int_0^L f(x) \cos \frac{n\pi}{L} x dx$$

(b) in a sine series

假設 $f(x) = -f(-x)$ for $-L < x < 0$, (假設 $f(x)$ 是一個 odd function)
interval 變為 $(-L, L)$

原本 sine series 公式

$$f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi}{p} x \quad b_n = \frac{2}{p} \int_0^p f(x) \sin \frac{n\pi}{p} x dx$$

現在只不過將 p 改成 L

$$f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi}{L} x \quad b_n = \frac{2}{L} \int_0^L f(x) \sin \frac{n\pi}{L} x dx$$

(c) in a Fourier series

interval 仍為 $(0, L)$

$$f(x) = f(x+L)$$

404

原本 Fourier series 公式

$$x \in [-p, p], f(x) = f(x+2p)$$

$$L = 2p$$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi}{p} x + b_n \sin \frac{n\pi}{p} x \right) \quad a_0 = \frac{1}{p} \int_{-p}^p f(x) dx$$

$$a_n = \frac{1}{p} \int_{-p}^p f(x) \cos \frac{n\pi}{p} x dx \quad b_n = \frac{1}{p} \int_{-p}^p f(x) \sin \frac{n\pi}{p} x dx$$

*4

Note! 不同一

不二

現在 (1) 將 interval $[-p, p]$ 換為 $[0, L]$, (2) 將 p 換為 $L/2$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{2n\pi}{L} x + b_n \sin \frac{2n\pi}{L} x \right) \quad a_0 = \frac{2}{L} \int_0^L f(x) dx$$

$$a_n = \frac{2}{L} \int_0^L f(x) \cos \frac{2n\pi}{L} x dx$$

$$b_n = \frac{2}{L} \int_0^L f(x) \sin \frac{2n\pi}{L} x dx$$


$$\text{Note: Since } f(x) \cos \frac{2n\pi}{L} x = f(x+L) \cos \frac{2n\pi(x+L)}{L}$$

$$\begin{aligned} & \int_{-L/2}^0 f(x) \cos \frac{2n\pi}{L} x dx \\ &= \int_{L/2}^L f(x) \cos \frac{2n\pi}{L} x dx \end{aligned}$$


$$a_n = \frac{2}{L} \int_{-L/2}^{L/2} f(x) \cos \frac{2n\pi}{L} x dx = \frac{2}{L} \int_0^L f(x) \cos \frac{2n\pi}{L} x dx$$

Example 3, $f(x) = x^2$, $0 < x < L$

將三個方法的結果畫成圖形




cosine series



sine series

from page 383

$$\begin{aligned}
 f_i(L) &= \frac{f(L^+) + f(L^-)}{2} \\
 &= \frac{f(-L^+) + f(L^-)}{2} \\
 &= \frac{-L^2 + L^2}{2} = 0
 \end{aligned}$$



Fourier series

$$\begin{aligned}
 f_1(L) &= \frac{f(L-) + f(L+)}{2} \\
 &= \frac{L^2 + f(0+)}{2} \\
 &= \frac{L^2}{2}
 \end{aligned}$$

11.3.6 Solving Particular Solutions (第四個方法)

$$a_n y^{(n)}(t) + a_{n-1} y^{(n-1)}(t) + \cdots + a_1 y'(t) + a_0 y(t) = f(t)$$

☆☆₂

$$\underline{f(t) = f(t + 2p)}$$

方法的限制

(註：以下的步驟不包含解 homogeneous solution
homogeneous solution 還是需要用 Section 4-3 的方法來解)

(Step 1) 將 $f(t)$ 表示成 Fourier series

$$f(t) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi}{p} t + b_n \sin \frac{n\pi}{p} t \right)$$

或 cosine series (當 $f(t)$ 為 even)

或 sine series (當 $f(t)$ 為 odd)

(Step 2) 假設 particular solution 的型態為

$$\underline{y_p(t) = A_0 + \sum_{n=1}^{\infty} \left(A_n \cos \frac{n\pi}{p} t + B_n \sin \frac{n\pi}{p} t \right)}$$
☆☆₂

(Step 3) 代回原式，比較係數，將 A_0, A_n, B_n 解出來

若所假設的 particular solution 和 homogeneous solution
有相同的地方，則要乘上 t

Example 4 (text page 441)

$$\text{auxiliary} = \frac{m^2}{16} + 4 = 0 \quad m^2 = -64 \\ m = \pm 8 \quad x_c(t) = C_1 \cos 8t + C_2 \sin 8t$$

$$\frac{1}{16} \frac{d^2x}{dt^2} + 4x = f(t)$$

$$f(t) = \pi t \quad \text{for } -1 < t < 1$$

$$f(t) = f(t-2) \quad P=1$$



(相關的物理定理見 Section 5-1)

Fourier sine series

(Step 1) 假設 $f(t) = \sum_{n=1}^{\infty} b_n \sin n\pi t$ (因為 $f(t)$ 是 odd) $f(t) = -f(-t)$

$$b_n = 2 \int_0^1 \pi t \sin(n\pi t) dt$$

$$= -2 \frac{t}{n} \cos(n\pi t) \Big|_0^1 + \int_0^1 \frac{2}{n} \cos(n\pi t) dt$$

$$= -2 \frac{1}{n} (-1)^n - 0 + \frac{2}{n^2 \pi} \sin(n\pi t) \Big|_0^1 = \frac{2}{n} (-1)^{n+1}$$

$$f(t) = \sum_{n=1}^{\infty} \frac{2}{n} (-1)^{n+1} \sin n\pi t$$

(Step 2) 假設 particular solution 為

$$x_p(t) = \sum_{n=1}^{\infty} (A_n \cos n\pi t + B_n \sin n\pi t) \quad (p = 1)$$

思考：為什麼這裡可以沒有常數項 A_0 ？

(Step 3) 將 $x_p(t)$ 和 Step 1 的結果代入 $\frac{1}{16} \frac{d^2x}{dt^2} + 4x = f(t)$

$$\sum_{n=1}^{\infty} \left(-\frac{1}{16} A_n n^2 \pi^2 \cos n\pi t - \frac{1}{16} B_n n^2 \pi^2 \sin n\pi t \right)$$

$$+ \sum_{n=1}^{\infty} (4A_n \cos n\pi t + 4B_n \sin n\pi t) = \sum_{n=1}^{\infty} \frac{2}{n} (-1)^{n+1} \sin n\pi t$$

$$-\frac{1}{16} A_n n^2 \pi^2 + 4A_n = 0 \quad \longrightarrow \quad A_n = 0$$

$$-\frac{1}{16} B_n n^2 \pi^2 + 4B_n = \frac{2}{n} (-1)^{n+1} \quad \longrightarrow \quad B_n = \frac{32(-1)^{n+1}}{n(64 - n^2 \pi^2)}$$

Therefore, the particular solution is:

$$x_p(t) = \sum_{n=1}^{\infty} \frac{32(-1)^{n-1}}{n(64 - n^2 \pi^2)} \sin n\pi t$$

General solution:

$$x(t) = c_1 \cos(8t) + c_2 \sin(8t) + \sum_{n=1}^{\infty} \frac{32(-1)^{n-1}}{n(64 - n^2\pi^2)} \sin n\pi t$$

注意：由於 $\frac{1}{16} \frac{d^2x}{dt^2} + 4x = f(t)$ 當中並沒有一次，三次，五次....微分項，所以 particular solution 不可能會有 cosine terms

所以，在 Step 2 當中，可以直接假設

$$x_p(t) = \sum_{n=1}^{\infty} B_n \sin n\pi t$$

11.3.7 Section 11.3 需要注意的地方

(1) 公式一些地方易記錯

for cosine series and sine series,

$$a_0 = \frac{2}{p} \int_0^p f(x) dx \quad a_n = \frac{2}{p} \int_0^p f(x) \cos \frac{n\pi}{p} x dx$$

$$b_n = \frac{2}{p} \int_0^p f(x) \sin \frac{n\pi}{p} x dx$$

(2) Fourier series 的 half-range extension 和 cosine series 及 sine series 不同

p is replaced by $L/2$, $[-p, p]$ is replaced by $[0, L]$

(3) Half range extension 和 solving particular solution 這兩個部分較複雜，需要特別注意，並且多練習例題

Exercise for Practice

Section 11-1 3, 5, 6, 8, 14, 17, 19, 20, 21, 22, 23

Section 11-2 2, 5, 9, 10, 12, 16, 19, 22, 23, 24

Section 11-3 14, 16, 18, 21, 22, 23, 28, 29, 33, 36, 37, 43, 46, 47a, 48a,
49, 52

Review 11 6, 12, 13, 14, 15, 17, 18